

The Petrochemistry and Origin of Precambrian Gneisses Around Madaka and Kwana-Bala Area, North Central Nigeria

Akintola.O. A^{1*} Adeleye. R.A²

*¹ Department of Geosciences, Nigerian Institute of Mining and Geosciences, Tudun Wada, Jos, Plateau State, Nigeria.

² Department of Geology, Afe Babalola University, Ado Ekiti, Ekiti State, Nigeria.

Corresponding Author: Akintola .O.
A

Article History

Received: 30/10/2024

Accepted: 05/11/2024

Published: 02/12/2024

Abstract: *This paper discusses the petrochemistry and origin of gneisses within basement complex of Nigeria around Madaka and Kwana-Bala area. The lithological units of rocks from Madaka and Kwana-Bala area comprises of gneisses (migmatitic, banded, granitic), kyanite-sillimanite schist, semi-pelitic schist, amphibolites, talcose rock, quartzite, phyllite, granodiorite, fine-medium-grained granites and porphyritic granite. Seven samples of gneisses were analyzed for major element oxides using (ICM-MS). The gneisses are often variable in colour which depending on the nature and proportion of the feldspar and the ferromagnesian minerals. The variation in textures of these rocks from study area is due to varying size and amount of the K-feldspar porphyroblasts couple with varying degree of foliation. In thin section, the characteristics granoblastic textures were observed in migmatitic and granitic gneisses. The mineral compositions of the gneisses rocks are variable, the common minerals assemblage includes: migmatitic gneiss quartz + plagioclase + microcline + biotite + chlorite + garnet, banded gneiss: quartz + plagioclase + muscovite + biotite + sphene + chlorite + garnet and granitic gneiss contain quartz + plagioclase + microcline + muscovite + biotite + chlorite + garnet + opaque minerals. The gneisses of the study area are product of reworked rocks during the Pan-African with a positive DF values, this suggests igneous sources for the protoliths of these rocks in the study area. The chemistry of the gneissic rock also show a variable proportion of Na₂O and K₂O with alkali-calcic affinity, which suggests, a closed system behaviour during metamorphism on one hand, and possibly a co-magmatic origin of the igneous protolith on the other hand.*

Keywords: Magmatic, Metamorphism, Petrochemistry, Precambrian, Protoliths, Tegna.

Introduction

The migmatite–gneiss of Nigeria basement complex is generally considered as the sensu stricto and it is the most widespread of the component units in the Precambrian basement [1, 2, 3], Figure 1. [4] revealed two generation of migmatitic-gneiss of different ages in the Northwestern basement blocks of Nigeria: The early migmatite type are differentiated in the field from the Pan-African Migmatite by complex fold structures with several generation and types of leucosomes which include granitic, aplitic and pegmatitic units, the Pan-African Migmatite show simple structures which include alternating schistose and tonalitic, granitic or pegmatitic layers. [4] Observed that the relationship between the gneisses and

the granitic gneiss suggests a common deformational episode. [5] Observed that the deformation and migmatitic processes that affected the basement tend to have dragged rocks of different lithologies, ages, and origin in parallelism such that differentiating these basement rocks into periodic episodal granitic additions becomes a problems. Geochronology of the gneiss complex from various parts of the Nigeria indicates that the rock has gone through at least four major tectonothermal events namely, the Liberian (2,700Ma, [5] Eburnean (2000Ma, [6], Kibarian (1,100Ma. [7] and Pan-African 600Ma [8], [9].




Figure 1: The geological map of Nigeria basement complex and other units of rocks [3].

Other reasonable age determinations are those by [10, 11, 12]. Previous works from Madaka and Kwana-Bala area includes petrographic descriptions of rocks [13, 14, 15, 16, 17, 18, 19, 20]. The most comprehensive study of any mineral group was done for the talc and amphiboles, their compositions were variables in Tegina area [15, 16, 19]. The geochemistry of rocks identified by [14] in Niger and Zaria provinces, have been studied [21, 22, 23,

24, 25, 26, 27]. Other works from Madaka and Kwana-Bala area includes the tectonic evolution of granites, amphibolites, and talcose rocks [6, 7, 15, 19]. This paper is therefore intended to discuss the petrochemistry of the gneisses, particularly in relation to their petro-tectonic significance in the Madaka and Kwana-Bala area Figure 2.




Figure 2: The location and accessibility map of the Madaka and Kwana-Bala area.

Materials and Methods

Seven (7) Fresh samples of gneisses were taken during the field work from Madaka and Kwana-Bala area with the aid of sledge hammer and chisel. The lithological units of rocks from Madaka and Kwana-Bala area were observed, described based on their mode of occurrence, macroscopic characteristics, structural patterns and field relation with adjacent rocks. Hand specimens were described based on the following macroscopic characteristics of the rocks. Samples taken from the field were prepared as thin section. The photomicrographs of obtained from thin sections of gneisses were produced in the petrographic Laboratory of Geology, Faculty of Sciences, Ahmadu Bello University, Zaria, Nigeria. Seven (7) selected samples of gneisses were analyzed for major element oxides at Activation Laboratories Limited (ACTLAB), analytical Laboratory Ancaster, Ontario, Canada.

Results

Field description and petrography of the gneisses

The basement rocks from Madaka and Kwana-Bala parts of Tegna Sheet 142 comprise of gneisses of varying proportion (migmatitic, bande and granitic), the metasediments includes kyanite-sillimanite schist, semi-pelitic schist, amphibolites, talcose rock, quartzite, and phyllite and granitic rocks consists of porphyritic granites, fine-medium-grained and granodiorite, Figure 3. The migmatitic gneisses around Gidian Zudangi and Gidan Biyaboki areas are cut by pegmatitic veins and have pygmatitic characteristic (Plate 1a). It is banded with leucocratic layer persistently thicker than the melanocratic layer. The migmatitic gneiss around Supana area were characterized with a variable development of veins and discontinuous streaks or augen of leucocratic layer most of which are parallel to the prevailing foliation (Plate 1b).




Figure 3: The geological map of the Madaka and Kwana-Bala area

Migmatitic gneiss around Kwaioki portrayed diktyonitic features (Plate 1a). Within migmatitic rock, layering is not conspicuous, the composition of quartz and plagioclase with minor amounts of biotite formed irregular, ovate or lenticular in shape. The biotites within this rock were surrounded by leucosome contents. The leucosome veins of the migmatitic gneiss fill shear, and locally tensile, fractures between enclaves of this rock Plate 1b. They are medium to coarse grained with the banding ranges from well banded to discontinuous in some outcrops. The northwestern and


southwestern part occurs as banded, granular in nature and consists of the plaeosome and neosome regions (Plate 1a). The plaeosome region (the gneissic) is the metamorphic host rock while the neosome consists of light grey medium grained granitic components which were introduced into the pre-existing parent rock that is the plaeosome (Plate 1b). Generally, the degree of migmatization increases in the vicinity of the principal granite and granitic gneiss masses.



A


Plate I Outcrops of migmatitic gneisses: (a) in form of low-lying along the northwestern part around Gidan Biyaboki with pygmatitic fold (Lat 10° 11' 47" N and Long 6° 21' 23" E), (B) showing diktyonitic features with enclaves surrounded by veins around Kwaioki area (Lat 10° 12' 02" N and Long 6° 29' 18" E)

The mafic minerals of the migmatitic gneiss (quartz and feldspar) from Madaka and Kwana-Bala area display discontinuous thin layers, lenses which were interleaved with layers rich in biotite as shown in (Plate IIb). The migmatitic gneiss of Madaka and Kwana-Bala area shows eudral to anhedral aggregates of dark and light minerals. The dark mineral is the biotite while the microcline, plagioclase and quartz are the light coloured minerals. The biotite minerals occurs along the cleavage, margins of the plagioclase and



B


as separate flakes. The biotite within the migmatitic gneiss tends to be distributed evenly and is oriented nearly parallel to the layering of the foliation planes. The quartz is colourless with low relief and had anhedral form. The quartz grains in this rock have irregular to lobate shapes and as a replacement toward the other minerals, the increasing alteration of the quartz within this rock unit became coarser, more irregular, and somewhat more abundant. The quantity of chlorite varies widely biotite occurred as light to deep brown felted mass due to Fe-rich materials with chlorite (Plate IIa-b). Plagioclase exists as tabular crystals. The migmatitic gneiss contain quartz 29%, plagioclase 18%, microcline 19%, biotite 24%, chlorite 4%, garnet 2% and opaque constitutes about 3% of the rock.



A


Plate V: The photomicrograph of migmatitic gneiss from Madaka and Kwana-Bala area under (a) plane polarized light (PPL) a mesosome region with the foliation defined by crystals, and annealed texture of unstrained quartz (b) crossed polarized light (XPL). Note contrast with a contrast in crystals shape between quartz (q) and feldspar (P)=Plagioclase (B)=Biotite, (Q)= quartz, M= Microcline C=Chlorite.

The banded gneiss in the study area occurs mostly as hills in the northwestern parts of the study area extending up to 300m around Gidan Sisi up to Gidan Usman. The banded gneiss occur as hilly and low lying outcrops, where they form contact with the schists. Contacts are generally gradational but sharp contacts are



B

commonly characterized with pre-tectonic dykes, pegmatite dyke and veins (Plates IIIa-b). The rock is dark to grey in colour. Individual bands of the banded gneisses vary greatly in thickness from a few millimetres to tens of centimetres Plate (IIIa). Texturally, it is fine to medium grained. Foliation within the bands, defined by aligned biotite and quartz, are almost always parallel to the banding. The proportion of neosome to paleosome varies in banded gneiss outcrop around Gidian Barau. Quartz, biotite and plagioclase contents of the paleosome also vary within wide limits. These variations are probably related to an original stratification. Paleosome is usually fine to medium grained, while the neosome is medium grained.



A




B


Plate III: Photographs of banded gneisses around (a) Gidan Sisi showing the extension of pre-tectonic dykes (Lat 10° 5' 47" N and Long 6° 16' 23" E), (b) Dutumi cut through by pegmatite dyke (Lat 10° 5' 31" N and Long 6° 17' 41"E),

The biotite contents in the banded gneisses occurs as pleochroic yellow-brown tabular aggregates. The quartz in the banded gneiss occurs as interstitial aggregates and as fine irregular grains characterized with micrographic intergrowths at the margin. The

feldspar (plagioclase) shows a deformed polysynthetic twinning free from zoning suggesting post crystal deformation in this rock (Plate IVb). The replacement of the plagioclase by sericite is also observed. The muscovite mineral occurs as coarse-grained, the mineral aggregates shows euhedral to subhedral in the banded gneiss. The mineral assemblage of the banded gneiss consists of quartz 17%, plagioclase 23%, muscovite 5%, biotite 43%, sphene 3%, chlorite 3%, garnet 3%, and opaque 3%.



A



B

Plate IV: The photomicrograph of banded gneiss from Madaka and Kwana-Bala area under (a) plane polarized light (PPL) with brown coloured biotite (b) crossed polarized light (XPL) showing biotite and plagioclase feldspar with deformed twin lamellae. B= Biotite, Q= Quartz, P= Plagioclase, M=Muscovite, C=Chlorite

The granitic gneiss is a highly foliated rock that is well-developed towards the Kwana-Bala border to the east and Godo to the west.

In the east of Makeri the rock covers an extensive area and rise into an oval shaped hill. Texturally, it is medium-grained, pale grey to white in colour. The granitic gneiss has a clear and sharp contact with the surrounding migmatitic gneiss (Plate Va). Structural features like faults and veins were mapped on the outcrop (Plate Vb).




A



B

Plate V: (a) The contact of granitic gneiss and migmatitic gneiss along Muazu-Wakayi road (Lat 11° 4' 46" N and Long 6° 19' 26" E), (b) fault and veins on the granitic gneiss along Godo road east of Supana (Latitude 10° 7' 22" N and Longitude 6° 22' 17" E).

The biotite occurs as flakes, fine to medium grained subhedral crystal. It is the most dominant mineral next to plagioclase in the rock. The biotite within the granitic gneiss occurred in two forms. One as light to deep brown felted mass of iron rich biotite without a well-defined crystal outline and the other as distinct green to pink



A

Plate VI: The photomicrograph of granitic gneiss from Madaka and Kwana-Bala area under (a) plane polarized

light (PPL) with foliation plane aligned with biotite (b) crossed polarized light (XPL) B= Biotite, Q= Quartz,


P= Plagioclase, H=hornblende, C= Chlorite, S=sphene.

Geochemistry

The major, trace and rare earth elements in gneisses from Madaka and Kwana-Bala areas were characterized using discriminants plots for quantitative comparison and genetic interpretation.

The major elements composition of the gneisses from Madaka and Kwaana-Bala area area indicates SiO₂ values of the gneisses range from 63.77- 69.9 wt%, average 65.66, Al₂O₃ ranges from 3.17- 6.88 wt%, average 15.9, Fe₂O₃(T) ranges from 13.46-17.3 wt%, average 4.84, CaO ranges from 2.15-3.64 wt%, average 3.01,

grains (PlateVIb). Quartz occurs in the form of large aggregate and lenses and generally have contacts with fragments of plagioclase. The plagioclase generally occurs as grey and platy crystals. Microcline crystals of varying sizes and shapes are common in the rock. The crystals show cross-hatch twinning. Accessory minerals include muscovite and garnet. The mineral assemblage of the granitic gneiss consists of quartz 30%, plagioclase 25%, microcline 10%, hornblende 5%, muscovite 3%, biotite 17%, chlorite 5%, garnet 3% and opaque 2%.



B

Na₂O ranges from 2.66-9.38 wt%, average 5.68 and L.O.I ranges from 2.57-4.0 wt%, average 2.68. In addition, the concentration of MnO, MgO, K₂O, TiO₂, P₂O₅ in the gneisses varies from 0.03-0.82 wt%, 0.011-0.822 wt%, 0.23-0.91 wt%, 0.01-0.75 wt%, 0.01-0.79 wt% respectively. The banded gneiss contains low P₂O₅ (0.01-0.04 wt%) compared to (0.34-0.68 wt%, 0.71-0.79 wt%) in granitic gneiss and migmatitic gneiss respectively. To determine the protolith of gneisses based on the amount of quartz and feldspatic composition, if the MgO< 6%, and SiO₂< 90% [28] with discriminant function (DF). [28] assigned positive and negative DF values to igneous and sedimentary rocks respectively using this general formular: Discriminant Function =10.44-0.21SiO₂ - 0.32Fe₂O₃T- 0.98MgO+0.55CaO+1.46Na₂O+0.54K₂O. Table (2) presents the DF of the gneisses obtained in the study area. The binary diagrams of Al₂O₃ versus MgO, after [29] (Figure 4) The variation diagram FeO/MgO versus FeO/MgO of the gneiss plots within high pressure metamorphism (Figure 5).

Table 1: Major elements composition and some ratios of gneisses in the study area (values in wt%).

	Granitic gneiss S ₂₅	Granitic gneiss S ₂₆	Granitic gneiss S ₂₇	Banded gneiss S ₂₈	Banded gneiss S ₂₉	Migmatitic gneiss S ₃₀	Migmatitic gneiss S ₃₁	Total	Ran Min	- ge Max	Average
Code											
Oxides											
SiO ₂	63.77	65.19	65.01	67.65	69.90	64.21	63.92	459.65	63.77	69.9	65.66
Al ₂ O ₃	15.90	17.30	17.11	14.41	13.46	16.63	16.98	111.79	13.46	17.3	15.9
Fe ₂ O ₃ (T)	4.10	3.41	3.17	4.84	5.22	6.21	6.88	33.83	3.17	6.88	4.84
MnO	0.41	0.82	0.73	0.07	0.03	0.58	0.71	3.35	0.03	0.82	0.48
MgO	0.01	0.09	0.011	0.18	0.11	0.8	0.822	2.02	0.011	0.82	0.29
CaO	3.19	3.22	2.15	3.64	2.33	3.34	3.2	21.07	2.15	3.64	3.01

Na ₂ O	9.38	6.11	7.40	5.99	5.11	2.66	3.11	39.76	2.66	9.38	5.68
K ₂ O	0.77	0.91	0.55	0.89	0.23	0.67	0.85	4.87	0.23	0.91	0.67
TiO ₂	0.01	0.27	0.09	0.01	0.27	0.11	0.75	1.51	0.01	0.75	0.22
P ₂ O ₅	0.34	0.68	0.71	0.01	0.04	0.79	0.71	3.28	0.01	0.79	0.47
L.O.I	2.12	2.0	3.07	2.67	3.30	4.0	1.57	18.73	1.57	4.00	2.68
Total	100 .0	100 .0	100.0	100.4	100.0	100 .0	100.0	100.36	100.0	100.4	100.1
FeO	3.69	3.07	2.85	4.35	4.69	5.58	6.19	4.35	2.81	6.19	4.35
FeO/MnO	9.00	3.74	3.90	62.1	157.3	9.62	8.72	254.38	3.74	157.3	36.34
FeO/MgO	369.0	34.11	385.0	24.17	42.64	6.98	7.75	769.65	6.98	369.0	109.95

Table 2. Discriminant function (DF) of the gneisses in the study area

Rock units	Discriminant function (DF) of the gneisses
Granitic gneiss S ₂₅	11.59
Granitic gneiss S ₂₆	7.88
Granitic gneiss S ₂₇	7.76
Banded gneiss S ₂₈	8.05
Banded gneiss S ₂₉	4.81
Migmatitic gneiss S ₃₀	0.27
Migmatitic gneiss S ₃₁	0.77




Figure 4: Bivariate Al₂O₃ versus MgO discrimination diagrams (after [29]) with line that separate paragneiss from orthogneiss for gneisses in the study area.




Figure 5: The binary diagram of FeO/MgO versus FeO/MnO in whole rock–comparison of different types of metamorphism (after [30]). Same key as in Figure 4)

Discussion

The gneisses (migmatitic gneiss, banded gneiss, granitic gneiss) of Madaka and Kwana-Bala area lies within the North central Nigerian Basement Complex. The rocks units within the study area comprise of the gneisses, metasediments and the granitoids (granodiorite, fine-medium-grained granites and porphyritic granite) as shown on the geological map of Madaka and Kwana-Bala area (Figure 4) with the exception of the minor rocks (the pegmatite and quartzite). The assemblages of these rocks show metamorphic imprints in the study area, indicative of various degree of deformation the rocks have suffered. The mineral composition of the gneisses in the study area is variable, the common assemblage being quartz + plagioclase +hornblende+ biotite and accessories like sphene and iron-a reflection of high amphibolites facies conditions during the D1 tectono-metamorphic episode. The crystallization of the major rock forming minerals in the gneisses, particularly in K-feldspar + plagioclase +biotite \pm muscovite controlled the assemblage. The occurrence of muscovite between plagioclase feldspar and quartz within the gneisses from Madaka and Kwana-Bala area, suggesting their late crystallization. The melting of the protolith of the gneisses occurred under hydrous conditions as indicated by the presence of hydrous minerals like biotite.

Field evidence from mineralogical descriptions of gneisses from Madaka and Kwana-Bala area indicates that the study area is a typical metamorphic zone where each rock units represents a state in the cycle of metamorphic change. The origin of the gneisses in Madaka and Kwana-Bala area has to be studied in terms of the general metamorphism that prevailed there and beyond. The gneisses are normally regarded as products of either injection of leucocratic, usually low-temperature and therefore more mobile, felsic minerals into pre-existing older bodies, or the segregation of such leucocratic components into separates bands that finally confer the lit-par-lit nature. In a regional metamorphic terranes, all these and more may take place which support the evolution models of the northern Nigerian Basement Complex by [23], [31]. The experiment of [32] suggests a closed-system derivation of gneiss (neosome) from clayey or smectitic family subjected to higher temperature of about 7000C at over 2 kb. In an open system, ingress by metasomatising fluids would progressively produce variants of metamorphites. Beginning with the lower-grade variants of what is evidently a metamorphic continuum. The assemblages of mineral, plagioclase-hornblende-biotite-garnet indicate that the metamorphic conditions were the lower grades of amphibolites facies in the ranges 4-8kb and 550-750oC (Winkler, 1979). It is importance to know if the gneisses around Madaka and Kwana-Bala area have acted as closed chemical systems or if some alterations process has changed the original chemistry of the rocks in the areas. The high concentrations of SiO₂ (63.77-69.9) in gneisses resulted from miscibility of silica contents of the rocks from Madaka and Kwana-Bala area (Table 1). The composition of CaO, NaO, K₂O, P₂O₅ and TiO₂ are strikingly lower to oxides of SiO₂, Al₂O₃ and Fe₂O₃ in the gneisses (Table 1). The lower values of these oxides were attributed to chemical reactions under oxidizing conditions of the source rocks in the Madaka and Kwana-Bala area. The gneisses from Madaka and Kwana-Bala area are product of reworking of the rocks during the Pan-African events as observed with the high contents of Na₂O and K₂O, predominantly of alkali-calcic affinity in nature, which can be

compared and related to rocks of Archaean migmatite gneisses [15], [27]. In line with [28] Table 2), the DF values of the gneissic rocks from Madaka and Kwana-Bala area including granitic gneiss (7.88-11.59), banded gneiss banded gneiss (4.81-8.05), migmatitic gneiss (0.27-0.77) are positive. Essentially, this suggests igneous sources for the gneisses in the area.

The binary plot of Al₂O₃ versus MgO diagram after [29] (Figure 4), shows that the gneisses falls in the orthogneiss field with one banded gneiss slightly lying in paragneiss field. The diagram FeO/MgO versus FeO/MgO after [30]. Figure 5 of the gneisses in the area plots within high pressure metamorphism [30, 31].

The origin of the gneisses from Madaka and Kwana-Bala area

The migmatitic gneiss from Madaka and Kwana-Bala area was a product of partial melting of the gneissic rock, leading to mineral segregation into mafic and felsic minerals in response to intense high grade metamorphism. They are presumed to be the oldest rock in the study area (Plate Ia-b). The banded gneiss from Madaka and Kwana-Bala area were form as a result of metamorphism of pelitic materials accumulated as older supracrustals. The banded gneiss from Madaka and Kwana-Bala area is characterized by alternation of mafic and felsic layers (Plate IIa-b) and the granitic gneiss from Madaka and Kwana-Bala area, composed of biotite, K-feldspars, quartz minerals, occurs in close association with the migmatitic gneiss. This suggests that the migmatitic rocks in the area were also intruded by granitic bodies probably during the early Kibaran event and were later metamorphosed together with the older supracrustals. This is evident also by cataclastic deformation which produced multiple features found in granitic gneiss. The veins occurring within the granitic gneiss are believed to have occurred as a result of syn deformational process during the last orogenic episode because it is concordant with the trend of the foliation (Plate IIIa-b). The gneisses of the study area are product of reworked rocks formed at the Pan-African event with a positive DF values, this suggests igneous sources for the protoliths of these rocks in the study area. This process was aided by partial melting of the protoliths.

Conclusion

Petrographic evidence of gneisses from Madaka and Kwana-Bala area show that, the Pan-African reworking of rocks led to re-crystallization of the minerals within the migmatite-gneiss complex (migmatitic, banded and granitic gneisses), with the lithology of rocks within the study area displaying variation of metamorphism facies (medium to upper amphibolites) in the area. The geochemistry data shows that the gneisses in the study area exhibit a high degree of geochemical variability in the major element oxides. The chemistry of the gneisses show a variable proportion of Na₂O and K₂O with alkali-calcic affinity, which suggest a closed system event during metamorphism of gneissic rocks on one hand and possibly a co-magmatic origin of the igneous protolith on the other hand.

Acknowledgments

Emeritus Prof. A.C Ajibade of Ajibade Garden's Ado-Ekiti is acknowledged by the first author for his helpful suggestions, comments and link up which significantly improved this work.

Conflict of interests.

No conflict of interests in this work

References

- McCurry, P. (1976). THE GEOLOGY OF THE PRECAMBRIAN TO LOWER PALAEOZOIC ROCKS OF NORTHERN NIGERIA. A REVIEW.
- Rahaman, M. A., Emofurieta, W. O., & Caen-Vachette, M. (1983). The potassic-granites of the Igbei area: further evidence of the polycyclic evolution of the Pan-African belt in southwestern Nigeria. *Precambrian Research*, 22(1-2), 75-92.
- Woakes, M., Rahaman, M. A., & Ajibade, A. C. (1987). Some metallogenetic features of the Nigerian basement. *Journal of African Earth Sciences* (1983), 6(5), 655-664.
- Ajibade, A. C., Woakes, M., & Rahaman, M. A. (1987). Proterozoic crustal development in the Pan-African Regime of Nigeria. *Proterozoic Lithospheric Evolution*, 17, 259-271.
- Oversby, V. M. (1975). Lead isotopic study of aplites from the Precambrian basement rocks near Ibadan, southwestern Nigeria. *Earth and Planetary Science Letters*, 27(2), 177-180.
- Kennedy Grant, N. (1970). Geochronology of Precambrian basement rocks from Ibadan, southwestern Nigeria. *Earth and Planetary Science Letters*, 10(1), 29-38.
- Grant, N. K., Rex, D. C., & Freeth, S. J. (1972). Potassium-argon ages and strontium isotope ratio measurements from volcanic rocks in northeastern Nigeria. *Contributions to Mineralogy and Petrology*, 35, 277-292.
- Ogezi, A. E. O. (1977). Geochemistry and geochronology of basement rocks from northwestern Nigeria (Doctoral dissertation, University of Leeds (Department of Earth Sciences)).
- Tubosun, I. A. (1983). GEOCHRONOLOGIE U/Pb DU SOCLE PRECAMBRIEN DU NIGERIA (Doctoral dissertation).
- Rahaman, M. A. (1988). Recent advances in the study of the basement complex of Nigeria. *Pre Cambrian geology of Nigeria*, 11-41.
- Dada, S. S. (1989). Evolution de la croute continentale au Nord Nigéria: apport de la géochimie, de la géochronologie U-Pb et des traceurs isotopiques Sr, Nd et Pb (Doctoral dissertation, Montpellier 2).
- Oyinloye, A. O. (2007). Geology and geochemistry of some crystalline basement rocks in Ilesha area, southwestern Nigeria: Implications on provenance and evolution. *Pakistan Journal of Scientific and Industrial Research*, 50.
- Truswell, J. F., & Truswell, J. F. (1960). A contribution to the study of older granites in the Nigerian Precambrian with special reference to the Tegina granite.
- Truswell, J. F., & Cope, R. N. (1963). The Geology of Parts of Niger and Zaria Provinces, Northern Nigeria: Explanation of 1: 250,000 Sheet No. 31. Federal Government of Nigeria.
- Elueze, A. A. (1981). Dynamic metamorphism and oxidation of amphibolites of Tegina area, northwest Nigeria. *Precambrian Research*, 14(3-4), 379-388.
- Amoka, I. S. (2000). Beneficiation of talcose rock from Kagara, North-central Nigeria. Unpublished Master of Technology Thesis, Federal University of Technology, Minna, Nigeria, 211.
- Omang, B. O., & Alabi, A. A. (2011). Geochemistry and mineralogical evaluation of quartzite bearing Kyanite in Kuta, Northwestern Nigeria. *Ethiopian Journal of Environmental Studies and Management*, 4(3), 79-83.
- Ihaza, C. A., Adekeye, J. I. D., & Omorinoye, O. A. (2014). Appraisal of the talc bodies of Kagara area, North-western Nigeria and their industrial potentials. *Centre point journal of science*, 20, 151-171.
- Akintola, O. A., Najime, T., & Abubakar, I. Y. (2017). Geochemistry and Origin of Talc Mineralization in Kagara Area (Sheet 142 SE and Part of Sheet 142 SW) North Central, Nigeria. *International Journal of Innovative Research and Development*, 6(12).
- Akintola, O. A., Najime, T., Adeleye R. A, Olaolun A. O, Ogunleye, P.O and Abubakar, I. Y. Petrography and mineralogy of manganese nodules around Madaka, Sheet 143, North Central, Nigeria, *Journal of Earth Science and Climatic Change*, 2022, 13(6), 25.
- Adeoye, A. O. The Composition and Origin of the Amphibolite's in Kumunu, North Central Nigeria.
- Ajibade, A. C. (1971). The geology of the country around Zungeru. Master of Science Thesis, University of Ibadan.
- Ajibade, A. C. (1976). PROVISIONAL CLASSIFICATION AND CORRELATION OF THE SCHIST BELTS IN NORTH-WESTERN NIGERIA.
- Grant, N. K. (1978). Structural distinction between a metasedimentary cover and an underlying basement in the 600-my-old Pan-African domain of northwestern Nigeria, West Africa. *Geological Society of America Bulletin*, 89(1), 50-58.
- Elueze, A. A. (1981). Geochemistry and Petrotectonic setting of metasedimentary rocks of the schist belt of Ilesa area, southwestern Nigeria. *Journal of Mining and Geology*, 18(1), 194-197.
- Ajibade, A. C., Fitches, W. R., & Wright, J. B. (1979). The Zungeru mylonites, Nigeria: Recognition of a major unit. *Rev de Geology. Geography*, 2, 359-363.

27. Dada, S. S. (2006). Proterozoic evolution of Nigeria. The basement complex of Nigeria and its mineral resources (A Tribute to Prof. MAO Rahaman). Akin Jinad and Co. Ibadan, 29-44.
28. Tourtelot, E. B. (1970). Selected annotated bibliography of minor-element content of marine black shales and related sedimentary rocks, 1930-65 (Vol. 1293). US Government Printing Office.
29. Meng, F., Nie, F., Xia, F., Yan, Z., Sun, D., Zhou, W., ... & Wang, Q. (2024). Geochemical characteristics and detrital zircon U-Pb ages of the Yimin Formation, Kelulun Depression, Hailar Basin and constraints on uranium mineralization. PloS one, 19(8), e0309433.
30. Chopin, C. (1984). Coesite and pure pyrope in high-grade blueschists of the Western Alps: a first record and some consequences. Contributions to Mineralogy and Petrology, 86, 107-118.
31. Ajibade, A. C., & Wright, J. B. (1989). The Togo-Benin-Nigeria Shield: evidence of crustal aggregation in the Pan-African belt. Tectonophysics, 165(1-4), 125-129.
32. Barker, A. J. (2013). Introduction to metamorphic textures and microstructures. Routledge.